

ISOLATION AND SCREENING OF ALKALI TOLERANT *TRICHODERMA SPP* AS BIOCONTROL AGENT FOR ALKALINE AGRICULTURAL SOIL

T. G. VIMALA KUMARI^{1*}, KAUSIK BASU¹, T. G. NITHYA², AJIT VARMA¹, AMIT C. KHARKWAL¹

¹Amity Institute of Microbial Technology (AIMT), Amity University Uttar Pradesh, Sector 125, Noida 201303, Uttar Pradesh, India,

²Department of Biotechnology, Faculty of Science and Humanities, SRM University Kattankulathur, India

Email: vimalasekaranprakash@yahoo.com

Received: 08 Sep 2014 Revised and Accepted: 05 Oct 2014

ABSTRACT

Objective: The main aim of the study was to screen twenty isolates of *Trichoderma spp* isolated from high alkaline agricultural soil and non agricultural soil and to test them *invitro* for their pH levels tolerance and biomass production.

Methods: All the twenty *Trichoderma spp* isolates were assessed for growth and biomass yield at different alkaline pH 9, 10, 11 on nutrient medium PCA and PCB. The growth were measured in terms of mycelial weight (cell biomass) and sporulation. The most tolerant isolates were further studied for their biocontrol activity against phytopathogen isolated from same location.

Results: Significant variations in the growth of the isolates were recorded with increasing pH. Different isolates of same species also varies in their growth and biomass yield at pH 7, 9, 10, and 11. The isolates were screened for highest growth at pH 11 in terms of biomass yield. 12 isolates showed prominent biomass yield at pH 10.0, but only 5 isolates were able to grow prominently at pH 11

Conclusion: The study concluded that *Trichoderma spp* isolated from alkali soil with a high pH of more than 11 was only able to grow under high pH range of 9, 10, 11 and also exhibited maximum antagonistic activity for the pathogens studied.

Keywords: *Trichoderma spp*, Alkaline pH, Tolerance, Biomass, Biocontrol.

INTRODUCTION

Biological control of a plant disease has always been thought to be the best remedial measure for disease management. It is an eco-friendly strategy of disease control and if the organism is self perpetuating, its repetitive applications are not required, thus reducing the cost of raising crops considerably. *Trichoderma spp* strains are of great importance as bio control agents, and should have better stress tolerance levels than the plant pathogens against which they are going to be applied for biological control under field condition [1]. *Trichoderma spp* is a genus of fungi that is present in all soils, where they are the most prevalent culturable fungi. Many species in this genus can be characterized as opportunistic avirulent plant symbionts[2]. *Trichoderma spp* have narrow and flexuous conidiophores and branches, with branches and phialides crowded, frequently paired, and seldom with more than three elements in a whorl[3].

The abiotic factors deteriorate the antagonistic properties of *Trichoderma spp*, against the phytopathogenic fungi [4]. Besides the effect of temperature, heavy metals, water relations, even the pesticides and pH have high influence on growth of phytopathogenic fungi as well as bio control agents. As in all microorganisms even in *Trichoderma spp*, the external factors modify its morphological characteristics as well as physiological functions [5].

Among these factors, pH is probably the most important environmental parameter affecting the mycoparasitic activities of *Trichoderma spp* strains [6]. A specific value of pH is required to note the maximum growth where these biocontrol agents can be multiplied and pathogen can be controlled. The studies on the variation of pH by different workers revealed that *Trichoderma spp* isolates showed optimum growth and sporulation rate at different pH values ranging from 2 to 7 [7].

Biocontrol agents need to be applied to rhizosphere for control of pathogen and enhancing plant growth. Studies suggested that these BCA failed to perform in soil with high alkaline pH. There is a serious problem in initial establishment of these *Trichoderma spp* isolates when applied in alkaline soil and is a notable problem emerging at a very high rate in present agricultural scenario.

The basic aim was to study indigenous bio control agent for sustainability in high alkaline pH. The present study has been initiated in consideration of developing a novel *Trichoderma spp* bioinoculants for application in alkaline soil for disease control. The alkali tolerant *Trichoderma spp* strain will imparts value to modern agriculture.

MATERIALS AND METHODS

Collection and isolation of the Fungus

The experimental material consisted of twenty isolates of *Trichoderma spp*, out of which 15 strains were isolated from agricultural soil sample collected from two district of Haryana and Gujarat-India and other 5 isolates from non agricultural soil. All the isolates were identified up to species level on the basis of phenotypic and morphological characteristics like colony colour and growth size and shape and branching of conidiophores, patterns and distribution of phialides and its shapes and conidia. The cultures were identified using the available literature [8,9,10, 11, 12].

Effect of pH on growth of *Trichoderma spp* isolates

Twenty *Trichoderma spp* isolates were assessed for growth and biomass yield at different alkaline pH 9, 10, 11 on nutrient medium PCA and PCB. The pH was adjusted accordingly. All the treatments were carried out in triplicates. The growth were measured in terms of mycelial weight (cell biomass) and sporulation. The data obtained were subjected to analysis of variance (ANNOVA). Mycelia biomass was considered as the most important determinants and significance of variance is presented in terms of 5% LOS.

Evaluation of Biocontrol properties

Fourteen phytopathogen were isolated from the same location and are known to cause minor or major disease symptoms in a range of plant species. Dual cultures were performed for determination of diffusible metabolites by *Trichoderma spp* isolate IPL/R &D/ VP/T2 and isolated phytopathogenic test fungus were grown on PDA plates. 5 mm disc of both the fungi (*Trichoderma spp*. and phytopathogenic test fungus) were placed on PDA plate about 2.0 – 2.5 cm away from each other. The plates were incubated at 25°C for one week.

Inhibition of the pathogenic fungal growth was measured. Control was taken without *Trichoderma spp* in the bottom plate. Observation was recorded after one week and percent inhibition was calculated. All the experiments were conducted in three replications.

RESULTS

Twenty isolates were obtained from the soil sample collected from two different states of Haryana and Gujarat -India and that involved both agricultural and non agricultural soil. The isolates were identified using various morphological keys available. The isolated *Trichoderma spp* species were differentiated on the basis of conidiophore and conidium morphology.

Isolation of *Trichoderma spp* isolates

A total of twenty isolates were screened and out of these, 15 isolates were from agricultural soil and 5 isolates were from non agricultural soil. The agricultural soil consists of varied crops like paddy, ground nut, cucumber, onion and so many.

The non agricultural soils were from areas adjacent to land with alkaline pH. The *Trichoderma spp* isolated from agricultural land are *Trichoderma sppviride*, *Trichoderma sppharzianum*, *Trichoderma sppasperellum*, *Trichoderma sppvirens* and *Trichoderma spplongibrachiatum*. *Trichoderma spp* obtained from non agricultural soils are *Trichoderma sppvirens*, *Trichoderma sppharzianum*, *Trichoderma sppkoningii*.

Effect of pH on growth of *Trichoderma spp* isolates

All the isolates were subjected to *invitro* study for alkali tolerance at pH 9,10,11,12 with pH 7 as control. Only five isolates showed a positive growth at pH 11.0.

Statistical Analysis

The data was subjected to analysis of variance (ANOVA) Table3:ANNOVA (Analysis of Variance) and significance of variance was presented at 5% level using IRRISTAT windows ver. 4.1.

Table 1: Isolated *Trichoderma spp* spp. from agricultural and non-agricultural soils

S. No.	Isolate No.	Organism
01	IPL/VP/01	<i>Trichoderma sppvirens</i>
02	IPL/VP/02	<i>Trichoderma sppviride</i>
03	IPL/VP/03	<i>Trichoderma sppvirens</i>
04	IPL/VP/04	<i>Trichoderma harzianum</i>
05	IPL/VP/05	<i>Trichoderma sppviride</i>
06	IPL/VP/06	<i>Trichoderma harzianum</i>
07	IPL/VP/07	<i>Trichoderma sppasperellum</i>
08	IPL/VP/08	<i>Trichoderma sppkoningii</i>
09	IPL/VP/09	<i>Trichoderma sppasperellum</i>
10	IPL/VP/10	<i>Trichoderma sppkoningii</i>
11	IPL/VP/11	<i>Trichoderma sppharzianum</i> ,
12	IPL/VP/12	<i>Trichoderma sppkoningii</i> .
13	IPL/VP/13	<i>Trichoderma sppvirens</i>
14	IPL/VP/14	<i>Trichoderma spplongibrachiatum</i>
15	IPL/VP/15	<i>Trichoderma sppvirens</i>
16	IPL/VP/16	<i>Trichoderma viride</i>
17	IPL/VP/17	<i>Trichoderma sppvirens</i>
18	IPL/VP/18	<i>Trichoderma harzianum</i>
19	IPL/VP/19	<i>Trichoderma harzianum</i>
20	IPL/VP/20	<i>Trichoderma harzianum</i>

Table 2: Effect of pH 7, 9, 10,11, 12 on mycelia biomass (g) of Twenty *Trichoderma spp* isolates

S. No.	Isolate No.	pH 7 Control in grams/100 ml of broth				pH 9.0				pH 10.0				pH 11.0			
		3 rd day	5 th day	7 th day	10 th day	3 rd day	5 th day	7 th day	10 th day	3 rd day	5 th day	7 th day	10 th day	3 rd day	5 th day	7 th day	10 th day
1	IPL/VP/01	0.7056	0.857	1.234	1.334	0.654	0.789	1.123	1.456	0.234	0.342	0.987	0.987	0.214	0.242	0.387	0.387
2	IPL/VP/02	0.9087	1.023	1.354	1.456	0.986	1.345	1.987	2.435	0.987	1.234	1.987	2.426	0.876	1.209	1.876	2.123
3	IPL/VP/03	0.876	0.945	1.357	1.396	0.834	0.945	1.342	1.301	0.345	0.345	0.879	0.873	0.112	0.213	0.479	0.579
4	IPL/VP/04	0.973	1.123	1.456	1.582	0.832	1.234	1.432	1.578	0.832	1.567	1.789	1.589	0.745	1.456	1.678	1.567
5	IPL/VP/05	0.543	0.786	0.932	1.153	0.456	0.687	0.823	1.023	0.234	0.345	0.945	1.008	0.132	0.123	0.564	0.987
6	IPL/VP/06	0.721	0.834	1.123	1.375	0.562	0.748	1.345	1.456	0.234	0.345	0.346	0.456	0.112	0.223	0.345	0.321
7	IPL/VP/07	0.675	0.723	0.983	0.985	0.543	0.653	0.897	0.927	0.321	0.345	0.876	0.867	0.223	0.245	0.576	0.687
8	IPL/VP/08	0.675	0.776	0.887	1.136	0.576	0.625	0.778	1.234	0.231	0.321	0.567	0.675	0.234	0.268	0.456	0.458
9	IPL/VP/09	0.765	0.908	1.123	1.235	0.723	0.823	1.205	1.298	0.823	0.856	1.605	1.698	0.786	0.869	1.505	1.678
10	IPL/VP/10	0.678	0.734	1.345	1.456	0.689	0.654	1.234	1.345	0.389	0.354	0.734	0.745	0.234	0.243	0.654	0.654
11	IPL/VP/11	0.556	0.886	1.586	1.589	0.435	0.713	1.432	1.123	0.224	0.223	0.334	0.567	0.112	0.119	0.245	0.345
12	IPL/VP/12	0.434	0.678	1.457	1.435	0.334	0.546	1.231	1.267	0.233	0.345	0.879	0.978	0.345	0.378	0.678	0.789
13	IPL/VP/13	0.786	0.978	1.345	1.457	0.345	0.567	1.210	1.100	0.145	0.456	0.710	0.900	0.104	0.234	0.564	0.700
14	IPL/VP/14	0.879	0.978	1.732	1.832	0.789	0.856	1.508	1.556	0.389	0.467	0.908	0.956	0.334	0.412	0.907	0.934
15	IPL/VP/15	0.765	0.9872	1.879	2.0321	0.556	0.872	1.259	1.523	0.456	0.572	1.005	1.023	0.123	0.145	0.605	0.403
16	IPL/VP/16	0.897	0.975	1.345	1.578	0.712	0.876	1.678	1.789	0.612	0.976	1.378	1.889	0.512	0.776	1.878	1.889
17	IPL/VP/17	0.432	0.6578	1.472	1.541	0.034	0.056	0.167	0.310	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.002
18	IPL/VP/18	0.669	0.789	0.856	0.893	0.567	0.589	0.786	0.987	0.267	0.489	0.786	0.887	0.112	0.438	0.656	0.778
19	IPL/VP/19	0.632	0.789	0.993	1.456	0.546	0.754	0.853	1.345	0.446	0.654	0.953	1.001	0.245	0.546	0.753	0.998
20	IPL/VP/20	0.856	0.9231	1.567	1.5890	0.856	0.765	1.456	1.345	0.231	0.564	1.045	1.231	0.112	0.345	1.123	1.200

A significant variation in biomass production was recorded among all isolates of *Trichoderma spp* species as well as different isolates of same species at all test pH values of 7, 9, 10 and 11 [Table 1]. Maximum number of isolates showed high biomass production at pH

7.00 followed by 9.00 minimum at pH 10.00only 5 isolates shows prominent growth at pH 11.00. The biomass production after 10 days i. e., at the end of the experiment ranged from 0.00 to 2.132 g in all treatments. With increasing incubation maximum isolates

showed a significant increase in biomass at all pH levels 7 – 9, but there was no significant increase in biomass at pH 10 and 11 with time. The biomass production of *T. viride*(IPL/VP/02, IPL/VP/16) was significantly higher than any other isolate at all pH levels 11 whereas *T. harzianum*(IPL/VP/04, IPL/VP/20); *T. asperellum* (IPL/VP/09) showed moderate biomass production. Minimum or no growth at pH 9, 10 was observed with *T. virens* (IPL/VP/17).

Evaluation of biocontrol properties

Statistical significance of antagonistic effect of isolated *Trichoderma sppviride*IPL/VP/02 was tested by 't' test (P = 0.05), against 14 isolated phyto pathogen and were expressed as mean ± SD of three experiments, and were subjected for analysis of variance. The dual culture study reveals that there is an antagonistic effect of the alkali

tolerant *Trichoderma sppviride* IPL/VP/02, though considerable variation in biocontrol properties with different pathogen.

DISCUSSION

A significant variation in the biomass of all the *Trichoderma spp* isolates was observed at different pH levels. The *Trichoderma spp* isolates from both agricultural soils and non agricultural soil showed their preference of pH 7 for optimal growth. Of all species, the maximum biomass was obtained for strains of *T. Viride* isolated from agricultural soil and also it was noticed that it utilized nutrients better than any other species at varied pH levels. *T. Viride* isolates also showed most inhibitory effect on the growth of phytopathogen isolated. Four other strains including strains of *T. harzianum* and *T. asperellum* shows prominent mycelia growth at pH level 10-11.

Table 3: Analysis of Data and significant variation

Sources of variation	Sum of Squares (SS)	Degree of Freedom (d. f.)	Mean sum of squares(MSS)	F _{calculated}
Between Groups	35.93	15	2.396	17.55
Within Groups	40.81	299	0.1365	
Total	76.74	314		




Fig. 1: Colony of *Trichoderma spp* isolates IPL/R&D/VP/T2, grown in different alkaline pH 7, 9, 10, 11.

Table 4: Percent inhibition of isolated phytopathogen by *Trichoderma sppviride*IPL/VP/02

S. No.	Pathogen	% inhibition in radial growth
01	Pathogen 1	50 ± 0.651
02	Pathogen 2	33.8 ± 1.21
03	Pathogen 3	64.4 ± 1.345
04	Pathogen 4	63.8 ± 0.987
05	Pathogen 5	38.2 ± 2.25
06	Pathogen 6	45.5 ± 1.36
07	Pathogen 7	23 ± 1.121
08	Pathogen 8	70.5 ± 1.201
09	Pathogen 9	44.2 ± 2.024
10	Pathogen 10	50 ± 1.354
11	Pathogen 11	35 ± 2.602
12	Pathogen 12	75.5 ± 2.121
13	Pathogen 13	60 ± 0.982
14	Pathogen 14	59 ± 3.121

Values are mean of three replicates + SD




Fig. 2: (1-14): Interaction of Isolated 14 pathogens with *Trichoderma spp* isolates IPL/R & D/ VP/T2.

But there is a considerable difference in the sporulation among the isolates. Only one *T. viride* isolates from agricultural soil sporulate with count as high as 2×10^8 was recorded. Some of the isolates showed biomass production at alkaline pH 9.00-11.00 but the sporulation was extremely low or no sporulation. Isolates from non agricultural soils preferred to grow at pH 7.0 – 10. This clearly indicated that the isolates obtained from agricultural soil have adapted themselves to different pH levels as the cultivation practices are known to affect the pH levels of agricultural soils which are reported to be alkaline [13]. It is different with the soils of non agricultural where there is no human intervention on the change of H⁺ ion concentration of soils. pH suitable for the *invitro* growth of the fungi was correlated with the pH of non agricultural forestsoil from where these fungi occur[14].

The present study also justify clearly that the isolates source plays an important role. The adaptation of *Trichoderma spp* isolates with varied pH clearly indicates that isolates from intense agricultural soils are tolerant to varied pH conditions. In contrast, isolates obtained from non- agricultural soils where there is no intervention of agricultural practices are not tolerant to varied pH levels. Hence, the present study recommend the use of these isolates as biocontrol agent (BCA) for obtaining better results in plant disease control. The isolate were tested for its biocontrol ability with 14 isolated phytopathogen and have shown significant results

CONCLUSION

The isolation source of *Trichoderma spp* isolates has an effect on the biomass production at different pH levels. The isolates from cultivated soils are more adaptive to the tested pH ranges than the isolates from virgin soils. Hence, the *Trichoderma* isolates isolated from high pH soils were able to grow and tolerate high pH and exhibited maximum antagonistic activity against the pathogens

isolated from the alkaline soils. Any ideal Biocontrol Agent has to survive and perform under stress conditions. Soil stress like high pH plays a major inhibitory role in preventing the most virulent culture to multiply and perform its activities. Hence stress tolerant *Trichoderma spp* especially high pH tolerant *Trichoderma spp* is required for the crop cultivation under alkaline soil conditions. This study hereby concludes that stress tolerant *Trichoderma spp* are present under naturally problematic soils and can be isolated successfully and used as biocontrol agent for those problematic soils.

CONFLICT OF INTERESTS

Declared None

REFERENCES

1. L Kredics, L Manczinger, Z Antal, Z Penzes, K Szekeres, F Kevel, et al. *In vitro* water activity and pH dependence of mycelial growth and extracellular enzyme activities of *Trichoderma spp* strains with biocontrol potential. J Applied Microbiol 2004;96:491-8.
2. Harman GE, Howell CR, Viterbo A, Chet I, Lorito M. "Trichoderma species—opportunistic avirulent plant symbionts". Nat Rev Microbiol 2004;2(1):43–56.
3. John. Bissett-A revision of the genus *Trichoderma* II. Infrageneric classification. Canadian J Bot 1991;69(11):2357-72.
4. J Dlużniewska. Reaction of fungi of *Trichoderma* genus to selected abiotic factors. Electronic J Agricultural University 2003;6:04.
5. Sarojini C Koll. Growth response of *Trichoderma* isolates against varying pH levels. Int J Environ Biol; 2012.
6. L. Kredics. Influence of environmental parameters on *Trichoderma spp* strains with biocontrol potential *Trichoderma spp* Strains with Biocontrol Potential. Food Technol Biotechnol 2003;41(1):37-42.

7. S Bandyopadhyay, S Jash, S Dutta. Effect of different pH and temperature levels on growth and sporulation of *Trichoderma* spp. *Environ Ecol* 2003;21:770-3.
8. J Bisset. A revision of the genus *Trichoderma* II. Infrageneric classification, Canadian. *J Bot* 1991a ;69:2357-72.
9. J Bisset. A revision of the genus *Trichoderma* III. Section *Pachybasium*, Canadian. *J Bot* 1991b ;69:2373-417.
10. J Bisset. A revision of the genus *Trichoderma* III. Additional notes on section *Longibrachiatum*, Canadian. *J Bot* 1991c;69:2418-20.
11. Nagamani IK Kunwar C. Manoharachary, Handbook of soil fungi, IK. International Private Limited: New Delhi; 2006. p. 409-25.
12. GJ Samuels, O Petrini, K Kuhls, E Lieckfeldt, CP Kubicek. The *Hypocrea schweinitzii* complex and *Trichoderma* sect. *Longibrachiatum*, *Studies Mycol* 1998;41:1-54.
13. Sarojini C Koll. Growth response of *Trichoderma* isolates against varying pH levels. *Int J Environ Biol*; 2012.
14. T Yamanaka. The effect of pH on the growth of saprophytic and ectomycorrhizal ammonia fungi *in vitro*. *Mycol* 2003;95:584-9.